

Ceritinib (Zykadia)

About This Drug

Ceritinib is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Pain in your abdomen
- Decreased appetite (decreased hunger)
- Tiredness
- Weight loss

Note: Each of the side effects above was reported in 25% or greater of patients treated with ceritinib. Not all possible side effects are included above.

Warnings and Precautions

- Severe and persistent diarrhea, pain in your abdomen, nausea and vomiting can occur.
- Changes in your liver function
- Inflammation (swelling) and/or scarring of the lungs. You may have a cough and/or trouble breathing.
- Abnormal heartbeat, electrocardiogram (EKG/ECG), and/or decrease in heart rate may occur which may be life-threatening
- Blood sugar levels may change
- Inflammation of your pancreas which can be life-threatening.

How to Take Your Medication

- Take this medicine by mouth with food.
- **Missed dose:** If you miss a dose, and it is less than 12 hours until your next dose, then skip the missed dose and go back to your normal schedule. If you miss a dose, and it is more than 12 hours until your next dose, take the missed dose. Do not take 2 doses at the same time or extra doses. Do not replace a vomited dose. If you vomit a dose or miss a dose, contact your doctor.

- **Handling:** Wash your hands after handling your medicine, your caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.
- **Storage:** Store this medicine in the original container at room temperature.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day. Be sure to include periods of rest between energy-draining activities.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- To help with weight loss, drink fluids that contribute calories (whole milk, juice, soft drinks, sweetened beverages, milkshakes, and nutritional supplements) instead of water.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.
- If you have diabetes, keep good control of your blood sugar level. Tell your nurse or your doctor if your glucose levels are higher or lower than normal.

Food and Drug Interactions

- Avoid grapefruit or grapefruit juice while taking this medicine as it may raise the levels of ceritinib in your body which could make side effects worse.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known drug interactions with ceritinib. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.



- Avoid the use of St. John's Wort while taking ceritinib as this may lower the levels of the drug in your body, which can make it less effective.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Feeling that your heart is beating in a fast or not normal way (palpitations)
- Pain in your chest or abdomen that does not go away
- Dry cough
- Wheezing and/or trouble breathing
- Feeling dizzy or lightheaded
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Tiredness that interferes with your daily activities
- Abnormal blood sugar
- Unusual thirst, passing urine often, headache, sweating, shakiness, irritability
- Signs of possible liver problems: dark urine, pale bowel movements, pain in your abdomen, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Severe pain in the abdomen that may spread to the back and worsen with eating
- If you think you may be pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for 6 months after treatment. Men with female partners of childbearing potential should use effective methods of birth control during your cancer treatment and for 3 months after your cancer treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for 2 weeks after treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised May 2021

