

## Dactinomycin (Cosmegen)

---

### About This Drug

Dactinomycin is used to treat cancer. It is given in the vein (IV).

### Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak (fatigue), and raise your risk of bleeding.
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Nausea and vomiting (throwing up)
- Decreased appetite (decreased hunger)
- Fever
- Tiredness
- Changes in your liver function
- Infection
- Difficulty swallowing
- Rash
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

**Note:** *Not all possible side effects are included above.*

### Warnings and Precautions

- Skin and tissue irritation including redness, pain, warmth, or swelling at the IV site if the drug leaks out of the vein and into nearby tissue. Very rarely it may cause tissue necrosis (death).
- Severe allergic skin reaction. You may develop blisters on your skin that are filled with fluid or a severe red rash all over your body that may be painful.
- Severe bone marrow suppression which can be life-threatening
- Changes in your liver function and blockage of small veins in the liver, which can cause liver failure and be life-threatening.
- Changes in your kidney function
- This drug may raise your risk of getting a second cancer such as leukemia.

- If you have received radiation treatments, your skin may become red after dactinomycin. This reaction is called “radiation recall.” Your body is recalling, or remembering, that it had radiation therapy.

**Note:** *Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team*

## Important Information

- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.
- Talk to your doctor before receiving any vaccinations during your treatment. Some vaccinations are not recommended while receiving dactinomycin.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have loose bowel movements, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or less these symptoms.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.



- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Avoid coloring your hair.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse. You can also call the American Cancer Society at 800-ACS-2345 to find out information about the “Look Good, Feel Better” program close to where you live. It is a free program where women getting chemotherapy can learn about wigs, turbans and scarves as well as makeup techniques and skin and nail care.
- If you received radiation, and your skin becomes red or irritated again, follow the same care instructions you did during radiation treatment. Be sure to tell the nurse or doctor administering your chemotherapy about your skin changes.

## Food and Drug Interactions

- There are no known interactions of dactinomycin with food.
- This drug may interact with other medicines. Tell your doctor and pharmacist about all the medicines and dietary supplements (vitamins, minerals, herbs and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

## When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Flu-like symptoms: fever, headache, muscle and joint aches, and fatigue (low energy, feeling weak)
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your mouth or throat that makes it hard to eat or drink
- Signs of possible liver problems: dark urine, pale bowel movements, bad stomach pain, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- Decreased urine, or very dark urine
- While you are getting this drug, please tell your nurse right away if you have any pain, redness, or swelling at the site of the IV infusion.
- A new rash or a rash that is not relieved by prescribed medicines
- Lasting loss of appetite or rapid weight loss of five pounds in a week



- If you think you may be pregnant or have impregnated your partner

## Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 6 months after your cancer treatment. Men with female partners of childbearing potential should use effective methods of birth control during your cancer treatment and for 3 months after your cancer treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for 14 days after treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Reviewed June 2020

