

Iron sucrose (Venofer)

About This Drug

Iron sucrose is used to treat anemia (low red blood cells) caused by low iron levels. It is given in the vein (IV).

Possible Side Effects

- Headache
- Feeling dizzy
- Pain in your chest
- Low blood pressure
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Itching
- Pain in your arm and/or leg
- Back pain
- Pain in your joints
- Muscle cramps
- Injection site reaction -you may get a rash, swelling or bruising or your skin may get red, warm, itchy, or painful at the site of your injection or infusion.
- Swelling in your legs, ankles, and/or feet

Note: Each of the side effects above was reported in 2% or greater of patients treated with iron sucrose. Not all possible side effects are included above.

Warnings and Precautions

- Allergic reactions, including anaphylaxis are rare but may happen in some patients and can be life threatening. Signs of allergic reaction to this drug may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this drug. You should get urgent medical treatment.**
- Severe low blood pressure

- Iron overload (too much iron in your body)

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

Treating Side Effects

- If you are dizzy, get up slowly after sitting or lying.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Moisturize your skin several times a day.
- Avoid sun exposure and apply sunscreen routinely when outdoors.
- While you are getting this drug, please tell your nurse right away if you get a rash, swelling or bruising or your skin gets red, warm, itchy, or painful at the site of your infusion or injection.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.

Food and Drug Interactions

- There are no known interactions of iron sucrose with food.
- This drug may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- A headache that does not away
- Feeling dizzy or lightheaded
- Tiredness that interferes with your daily activities
- Chest pain
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day



- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Itching that is bothersome
- Weight gain of 5 pounds in one week (fluid retention)
- Swelling in your legs, ankles, and/or feet
- Pain that does not go away or is not relieved by prescribed medicine
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- If you think you may be pregnant

Reproduction Warnings

- **Pregnancy warning:** It is not known if this drug may harm an unborn child. For this reason, be sure to talk with your doctor if you are pregnant or planning to become pregnant while receiving this drug. Let your doctor know right away if you think you may be pregnant.
- **Breastfeeding warning:** This drug passes into breast milk. For this reason, women should talk to their doctor about the risks and benefits of breastfeeding during treatment with this drug because this drug may enter the breast milk and cause diarrhea and constipation to a breastfeeding baby.
- **Fertility warning:** Human fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children.

Revised January 2021

