

Ixabepilone (Ixempra)

About This Drug

Ixabepilone is used to treat cancer. It is given in the vein (IV).

Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak (fatigue), and raise your risk of bleeding.
- Nausea and vomiting (throwing up)
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Diarrhea (loose bowel movements)
- Tiredness and weakness
- Joint, bone and muscle pain
- Effects on the nerves are called peripheral neuropathy. You may feel numbness, tingling, or pain in your hands and feet. It may be hard for you to button your clothes, open jars, or walk as usual. The effect on the nerves may get worse with more doses of the drug. These effects get better in some people after the drug is stopped but it does not get better in all people.
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: Each of the side effects above was reported in 20% or greater of patients treated with ixabepilone. Not all possible side effects are included above. Your side effects may differ if given in combination with capecitabine.

Warnings and Precautions

- Severe bone marrow suppression, which can be life-threatening
- Severe peripheral neuropathy
- Allergic reactions, including anaphylaxis are rare but may happen in some patients. Signs of allergic reaction to this drug may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this drug. You should get urgent medical treatment.**
- Changes in your liver function, which can cause liver failure

- Changes in the tissue of the heart and heart attack. Some changes may happen that can cause your heart to have less ability to pump blood.
- This drug contains dehydrated alcohol USP and may affect your central nervous system. The central nervous system is made up of your brain and spinal cord. You may feel dizzy and very sleepy.

Note: *Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.*

Important Information

- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.
- This drug may impair your ability to drive or use machinery. Use caution and tell your nurse or doctor if you feel dizzy or very sleepy.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have loose bowel movements, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.



- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.
- Get regular exercise. If you feel too tired to exercise vigorously, try taking a short walk.
- If you have numbness and tingling in your hands and feet, be careful when cooking, walking, and handling sharp objects and hot liquids.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Avoid coloring your hair.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse. You can also call the American Cancer Society at 800-ACS-2345 to find out information about the “Look Good, Feel Better” program close to where you live. It is a free program where women getting chemotherapy can learn about wigs, turbans and scarves as well as makeup techniques and skin and nail care.

Food and Drug Interactions

- Avoid grapefruit or grapefruit juice while taking this medicine. Grapefruit and grapefruit juice may raise the levels of ixabepilone in your body. This could make side effects worse.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs and others) you are taking before starting this medicine as there are known drug interactions with ixabepilone. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- Avoid the use of St. John’s Wort while taking ixabepilone as this may lower the levels of the drug in your body, which can make it less effective.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness or extreme weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Trouble breathing
- Feeling that your heart is beating in a fast or not normal way (palpitations)
- Pain in your chest
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Pain in your mouth or throat that makes it hard to eat or drink
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Pain that does not go away, or is not relieved by prescribed medicines



- Numbness, tingling, or pain in your hands and feet
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- Signs of possible liver problems: dark urine, pale bowel movements, bad stomach pain, feeling very tired and weak, unusual itching, or yellowing of the eyes or skin
- If you think you may be pregnant

Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment. Let your doctor know right away if you think you may be pregnant.
- **Breastfeeding warning:** It is not known if this drug passes into breast milk. For this reason, women should talk to their doctor about the risks and benefits of breastfeeding during treatment with this drug because this drug may enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Human fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Reviewed July 2020

