

UPMC | HILLMAN CANCER CENTER

Levoleucovorin (Fusilev®)

About This Drug

Levoleucovorin is similar to a vitamin. It is used in combination with other cancer medications such as 5-fluorouracil and methotrexate. Levoleucovorin helps 5-fluorouracil kill the cancer cells. It also helps to reduce the side effects of methotrexate. Levoleucovorin is given in the vein (IV).

Possible Side Effects

- Nausea and vomiting (throwing up)
- Soreness of the mouth and throat. You may have red areas, white patches, or sores in your mouth that hurt.
- Diarrhea (loose bowel movements)
- Tiredness, weakness
- Decreased appetite (decreased hunger)
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.
- Inflammation of the skin/rash

Note: Each of the side effects above was reported in 19% or greater of patients treated with levoleucovorin. Not all possible side effects are included above.

Treating Side Effects

- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have loose bowel movements, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help or stop lessen these symptoms.
- If you get diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water

or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.

- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- To help with decreased appetite, eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste.
- Daily exercise, with your doctor's approval, may increase your appetite.
- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Avoid coloring your hair.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse. You can also call the American Cancer Society at 800-ACS-2345 to find out information about the "Look Good, Feel Better" program close to where you live. It is a free program where women getting chemotherapy can learn about wigs, turbans and scarves as well as makeup techniques and skin and nail care.
- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.

Food and Drug Interactions

- There are no known interactions of levoleucovorin with food.
- This drug may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.5 F (38 C) or above
- Chills
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Pain in your mouth or throat that makes it hard to eat or drink
- New rash and/or itching
- Rash that is not relieved by prescribed medicines

- Extreme weakness that interferes with normal activities
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way
- If you think you are pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** It is not known if this drug may harm an unborn child. For this reason, be sure to talk with your doctor if you are pregnant or planning to become pregnant while receiving this drug. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** It is not known if this drug passes into breast milk. For this reason, women should talk to their doctor about the risks and benefits of breastfeeding during treatment with this drug because this drug may enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Human fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Revised October 2018