

## Liposomal Irinotecan (Onivyde)

---

### About This Drug

Liposomal irinotecan is used to treat cancer. It is given in the vein (IV).

### Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak (fatigue), and raise your risk of bleeding.
- Tiredness and weakness
- Fever
- Soreness of the mouth and throat. You may have red areas, white patches, or sores that hurt.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Decreased appetite (decreased hunger)

**Note:** Each of the side effects above was reported in 20% or greater of patients treated with liposomal irinotecan. Not all possible side effects are included above.

### Warnings and Precautions

- Severe diarrhea, which can be life-threatening
- Fever and severe decrease in white blood cells which puts you at an increased risk of infections which can be life-threatening
- Allergic reactions, including anaphylaxis are rare but may happen in some patients. Signs of allergic reaction to this drug may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this drug. You should get urgent medical treatment.**
- Inflammation (swelling) of the lungs, which can be life-threatening. You may have a dry cough or trouble breathing.

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

## Important Information

- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Mouth care is very important. Your mouth care should consist of routine, gentle cleaning of your teeth or dentures and rinsing your mouth with a mixture of 1/2 teaspoon of salt in 8 ounces of water or 1/2 teaspoon of baking soda in 8 ounces of water. This should be done at least after each meal and at bedtime.
- If you have mouth sores, avoid mouthwash that has alcohol. Also avoid alcohol and smoking because they can bother your mouth and throat.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have loose bowel movements, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.



## Food and Drug Interactions

- This drug may interact with grapefruit and grapefruit juice. Talk to your doctor as this could make side effects worse.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs and others) you are taking before starting this medicine as there are known drug interactions with liposomal irinotecan. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- Avoid the use of St. John's Wort while taking liposomal irinotecan as this may lower the levels of the drug in your body, which can make it less effective.

## When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Tiredness and weakness that interferes with your daily activities
- Pain in your chest
- Dry cough
- Trouble breathing
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Pain in your mouth or throat that makes it hard to eat or drink
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**
- If you think you may be pregnant or may have impregnated your partner

## Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for 1 month after treatment. Men with female partners of childbearing potential should use effective methods of birth control during your cancer treatment and for 4 months after your cancer treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for 1 month after treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.



- **Fertility warning:** Human fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Reviewed July 2020

