

## Pertuzumab (Perjeta)

---

### About This Drug

Pertuzumab is used to treat cancer. It is given in the vein (IV).

### Possible Side Effects

- Bone marrow suppression. This is a decrease in the number of white blood cells, red blood cells, and platelets. This may raise your risk of infection, make you tired and weak (fatigue), and raise your risk of bleeding.
- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Not able to move bowels (constipation)
- Tiredness
- Headache
- Muscle pain/aching
- Effects on the nerves are called peripheral neuropathy. You may feel numbness, tingling, or pain in your hands and feet. It may be hard for you to button your clothes, open jars, or walk as usual. The effect on the nerves may get worse with more doses of the drug. These effects get better in some people after the drug is stopped but it does not get better in all people.
- Rash
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

**Note:** Each of the side effects above was reported in 30% or greater of patients treated with pertuzumab. Not all possible side effects are included above.

### Warnings and Precautions

- Congestive heart failure – your heart has less ability to pump blood properly. You may be short of breath. Your arms, hands, legs and feet may swell.
- Allergic reactions, including anaphylaxis are rare but may happen in some patients. Signs of allergic reaction to this drug may be swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, do not take another dose of this drug. You should get urgent medical treatment.**

- While you are getting this drug in your vein (IV), you may have a reaction to the drug. Sometimes you may be given medication to stop or lessen these side effects. Your nurse will check you closely for these signs: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. **These reactions may happen after your infusion. If this happens, call 911 for emergency care.**

**Note:** Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

## Important Information

- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.

## Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- To decrease the risk of infection, wash your hands regularly.
- Avoid close contact with people who have a cold, the flu, or other infections.
- Take your temperature as your doctor or nurse tells you, and whenever you feel like you may have a fever.
- To help decrease the risk of bleeding, use a soft toothbrush. Check with your nurse before using dental floss.
- Be very careful when using knives or tools.
- Use an electric shaver instead of a razor.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature. Ask your nurse or doctor about other helpful tips and medicine that is available to help stop or lessen these symptoms.
- If you throw up or have loose bowel movements, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea or constipation.
- If you are not able to move your bowels, check with your doctor or nurse before you use enemas, laxatives, or suppositories.
- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.
- If you have numbness and tingling in your hands and feet, be careful when cooking, walking, and handling sharp objects and hot liquids.
- Infusion reactions may occur after your infusion. **If this happens, call 911 for emergency care.**
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day.



- Avoid rubbing your scalp, pat your hair or scalp dry.
- Avoid coloring your hair.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse. You can also call the American Cancer Society at 800-ACS-2345 to find out information about the “Look Good, Feel Better” program close to where you live. It is a free program where women getting chemotherapy can learn about wigs, turbans and scarves as well as makeup techniques and skin and nail care.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.
- Get regular exercise. If you feel too tired to exercise vigorously, try taking a short walk.

## Food and Drug Interactions

- There are no known interactions of pertuzumab with food.
- This drug may interact with other medicines. Tell your doctor and pharmacist about all the prescription and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs and others) that you are taking at this time. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.

## When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- Fever of 100.4° F (38° C) or higher
- Chills
- Trouble breathing
- Tiredness or weakness that interferes with your daily activities
- Feeling dizzy or lightheaded
- Easy bleeding or bruising
- Swelling of arms, legs, ankles, or feet
- Weight gain of 5 pounds in one week (fluid retention)
- Headache that does not go away
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- No bowel movement in 3 days or when you feel uncomfortable
- A new rash or a rash that is not relieved by prescribed medicines
- Numbness, tingling, or pain in your hands and feet
- Signs of allergic reaction: swelling of the face, feeling like your tongue or throat are swelling, trouble breathing, rash, itching, fever, chills, feeling dizzy, and/or feeling that your heart is beating in a fast or not normal way. **If this happens, call 911 for emergency care.**



- Signs of infusion reaction: fever or shaking chills, flushing, facial swelling, feeling dizzy, headache, trouble breathing, rash, itching, chest tightness, or chest pain. **If this happens, call 911 for emergency care.**
- If you think you may be pregnant

## Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for 7 months after treatment. Let your doctor know right away if you think you may be pregnant.
- **Breastfeeding warning:** It is not known if this drug passes into breast milk. For this reason, women should not breastfeed during treatment and for 7 months after treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** Human fertility studies have not been done with this drug. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm or egg banking.

Reviewed July 2020

