

Ripretinib (Qinlock)

About This Drug

Ripretinib is used to treat cancer. It is given orally (by mouth).

Possible Side Effects

- Nausea and vomiting (throwing up)
- Diarrhea (loose bowel movements)
- Constipation (not able to move bowels)
- Pain in your abdomen
- Tiredness
- Decreased appetite (decreased hunger)
- Muscle pain
- Hand-foot syndrome. The palms of your hands or soles of your feet may tingle, become numb, painful, swollen, or red.
- Hair loss. Hair loss is often temporary, although with certain medicine, hair loss can sometimes be permanent. Hair loss may happen suddenly or gradually. If you lose hair, you may lose it from your head, face, armpits, pubic area, chest, and/or legs. You may also notice your hair getting thin.

Note: Each of the side effects above was reported in 20% or greater of patients treated with ripretinib. Not all possible side effects are included above.

Warnings and Precautions

- Severe hand-foot syndrome
- This drug may raise your risk of getting new skin cancers.
- High blood pressure, which can be severe
- Congestive heart failure – your heart is not pumping blood as well as it should be, and fluid can build up in your body.
- Slow wound healing

Note: Some of the side effects above are very rare. If you have concerns and/or questions, please discuss them with your medical team.

Important Information

- Ripretinib may cause slow wound healing. It should not be given within at least 1 week of surgery and for at least 2 weeks after major surgical procedures and until the wound is fully healed. If you must have emergency surgery or have an accident that results in a wound, tell the doctor that you are on ripretinib.

How to Take Your Medication

- Swallow the medicine whole with or without food as prescribed. Take this medicine at the same time each day it is scheduled.
- **Missed dose:** If you miss a dose by more than 8 hours, then skip the missed dose and go back to your normal schedule. If it has been less than 8 hours since your missed dose, then take the missed dose. Do not take 2 doses at the same time or extra doses. Do not replace a vomited dose. If you vomit a dose or miss a dose, contact your doctor.
- **Handling:** Wash your hands after handling your medicine. Caretakers should not handle your medicine with bare hands and should wear latex gloves.
- This drug may be present in the saliva, tears, sweat, urine, stool, vomit, semen, and vaginal secretions. Talk to your doctor and/or your nurse about the necessary precautions to take during this time.
- **Storage:** Store this medicine in the original container at room temperature with the desiccant to protect from moisture and light. Replace cap securely each time after opening. Do not remove desiccant.
- **Disposal of unused medicine:** Do not flush any expired and/or unused medicine down the toilet or drain unless you are specifically instructed to do so on the medication label. Some facilities have take-back programs and/or other options. If you do not have a take-back program in your area, then please discuss with your nurse or your doctor how to dispose of unused medicine.

Treating Side Effects

- Manage tiredness by pacing your activities for the day.
- Be sure to include periods of rest between energy-draining activities.
- Get regular exercise. If you feel too tired to exercise vigorously, try taking a short walk.
- Drink plenty of fluids (a minimum of eight glasses per day is recommended).
- If you throw up or have diarrhea, you should drink more fluids so that you do not become dehydrated (lack of water in the body from losing too much fluid).
- If you have diarrhea, eat low-fiber foods that are high in protein and calories and avoid foods that can irritate your digestive tracts or lead to cramping.
- Ask your nurse or doctor about medicine that can lessen or stop your diarrhea, constipation and/or nausea.
- If you are not able to move your bowels, check with your doctor or nurse before you use enemas, laxatives, or suppositories.
- To help with nausea and vomiting, eat small, frequent meals instead of three large meals a day. Choose foods and drinks that are at room temperature.



- To help with decreased appetite, eat small, frequent meals. Eat foods high in calories and protein, such as meat, poultry, fish, dry beans, tofu, eggs, nuts, milk, yogurt, cheese, ice cream, pudding, and nutritional supplements.
- Consider using sauces and spices to increase taste. Daily exercise, with your doctor's approval, may increase your appetite.
- Avoid sun exposure and apply sunscreen routinely when outdoors.
- If you get a rash do not put anything on it unless your doctor or nurse says you may. Keep the area around the rash clean and dry. Ask your doctor for medicine if your rash bothers you.
- To help with hair loss, wash with a mild shampoo and avoid washing your hair every day. Avoid coloring your hair.
- Avoid rubbing your scalp, pat your hair or scalp dry.
- Limit your use of hair spray, electric curlers, blow dryers, and curling irons.
- If you are interested in getting a wig, talk to your nurse and they can help you get in touch with programs in your local area.
- Keeping your pain under control is important to your well-being. Please tell your doctor or nurse if you are experiencing pain.

Food and Drug Interactions

- This drug may interact with grapefruit and grapefruit juice. Talk to your doctor as this could make side effects worse.
- Check with your doctor or pharmacist about all other prescription medicines and over-the-counter medicines and dietary supplements (vitamins, minerals, herbs, and others) you are taking before starting this medicine as there are known drug interactions with ripretinib. Also, check with your doctor or pharmacist before starting any new prescription or over-the-counter medicines, or dietary supplements to make sure that there are no interactions.
- Avoid the use of St. John's Wort while taking ripretinib as this may lower the levels of the drug in your body, which can make it less effective.

When to Call the Doctor

Call your doctor or nurse if you have any of these symptoms and/or any new or unusual symptoms:

- A headache that does not go away
- Blurred vision or other changes in eyesight
- Trouble breathing
- Feeling that your heart is beating fast or in a not normal way (palpitations)
- Tiredness and/ or extreme weakness that interferes with your daily activities
- Nausea that stops you from eating or drinking and/or is not relieved by prescribed medicines
- Throwing up more than 3 times a day
- Diarrhea, 4 times in one day or diarrhea with lack of strength or a feeling of being dizzy
- No bowel movement in 3 days or when you feel uncomfortable
- Pain in your abdomen that does not go away



- Muscle pain that does not go away, or is not relieved by prescribed medicines
- Lasting loss of appetite or rapid weight loss of five pounds in a week
- Swelling of the legs, feet, or ankles
- Weight gain of 5 pounds in one week (fluid retention)
- Painful, red, or swollen areas on your hands or feet
- Numbness and/or tingling of your hands and/or feet
- New skin sore, or reddish bump that bleeds or does not heal
- New wart
- Change in size or color of a mole
- If you think you may be pregnant or may have impregnated your partner

Reproduction Warnings

- **Pregnancy warning:** This drug can have harmful effects on the unborn baby. Women of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 1 week after treatment. Men with female partners of childbearing potential should use effective methods of birth control during your cancer treatment and for at least 1 week after your cancer treatment. Let your doctor know right away if you think you may be pregnant or may have impregnated your partner.
- **Breastfeeding warning:** Women should not breastfeed during treatment and for at least 1 week after treatment because this drug could enter the breast milk and cause harm to a breastfeeding baby.
- **Fertility warning:** In men, this drug may affect your ability to have children in the future. Talk with your doctor or nurse if you plan to have children. Ask for information on sperm banking.

Revised April 2021

