



COVID-19 Vaccination for Ages 18 and Older

Last Updated: 6/23/2022

Primary Series: The vaccination(s) needed to give the first basic immunity to a virus.

Booster Dose: The dose(s) used to remind a healthy immune system to remain prepared.

Additional Dose: May improve the chance of a response by a weakened immune system.


Talk with your doctor about when to get vaccinated if recently exposed or infected, if you have concerns about potential side effects, and about which vaccine is right for you.



Highly recommended - get it as soon as you are eligible



Suggested - may get to keep your immune system prepared



* Recommendation is to receive your 2nd dose closer to 8 weeks after your 1st dose if you are:

- Male aged 12-39 years to lower the rare risk of heart inflammation
- Aged 65 or older to increase your immune response to the vaccine

Recommendation is to receive your 2nd dose closer to 3 to 4 weeks after your 1st dose if:

- You are at risk for severe illness or at high risk for exposure to the virus

** Those ages 18-49 who received J&J for their primary series AND 1st booster dose can receive a 2nd booster dose of Pfizer or Moderna.

Continued >




COVID-19 Vaccination for Ages 18 and Older

For Those Who Are Immunocompromised

Highly recommended - get it as soon as you are eligible

Suggested - may get to keep your immune system prepared



Those who received a hematopoietic cell transplant, or a CAR-T cell treatment should discuss additional doses with their doctor.

*Talk with your doctor if you have any questions.

To learn more, please visit Vaccine.UPMC.com.

Continued >



COVID-19 Vaccination for Ages 6 Months to < 18 Years

Last Updated: 6/23/2022



Highly recommended - get it as soon as you are eligible



Suggested - may get to keep your immune system prepared

Primary Series: The vaccination(s) needed to give the first basic immunity to a virus.

Booster Dose: The dose(s) used to remind a healthy immune system to remain prepared.

Additional Dose: May improve the chance of a response by a weakened immune system.



Your Primary Series



Booster Dose

PFIZER 3 doses
ages 6 months to < 5 years

First 2 doses are 3-8 weeks apart. Third dose given 8 weeks or more after the 2nd dose.

Not authorized

PFIZER 2 doses
3-8 weeks apart*
ages 5 to < 18 years

5 months or more after 2nd dose

Which vaccine should I get? ✓ **Pfizer**

MODERNA 2 doses
4-8 weeks apart*
ages 6 months to < 6 years

Not authorized

MODERNA 2 doses
4-8 weeks apart*
ages 6 to < 18 years

Not authorized

*Recommendation is to receive your 2nd dose closer to 8 weeks after your 1st dose if you are:
• Male aged 12-39 years to lower the rare risk of heart inflammation
Recommendation is to receive your 2nd dose closer to 3 to 4 weeks after your 1st dose if:
• You are at risk for severe illness or at high risk for exposure to the virus



COVID-19 Vaccination for Ages 6 Months to < 18 Years

For Those Who Are Immunocompromised

- Highly recommended -** get it as soon as you are eligible
- Suggested -** may get to keep your immune system prepared

Your Primary Series	4th Dose	5th Dose
PFIZER 3 doses ages 6 months to < 5 years First 2 doses are 3 weeks apart. Third dose given 8 weeks or more after the 2nd dose.	Not authorized	Not authorized
PFIZER 3 doses ages 5 to < 12 years First 2 doses are 3 weeks apart. Third dose given 4 weeks or more after the 2nd dose.	3 months or more after 3rd dose	Not authorized
PFIZER 3 doses ages 12 to < 18 years First 2 doses are 3 weeks apart. Third dose given 4 weeks or more after the 2nd dose.	Which vaccine should I get? ✓ Pfizer	4 months or more after 4th dose Which vaccine should I get? ✓ Pfizer
MODERNA 3 doses ages 6 months to < 6 years First 2 doses are 4 weeks apart. Third dose given 4 weeks or more after the 2nd dose.	Not authorized	Not authorized
MODERNA 3 doses ages 6 to < 18 years First 2 doses are 4 weeks apart. Third dose given 4 weeks or more after the 2nd dose.	Not authorized	Not authorized

